

Medical Conditions & Drugs Associated with IIH

Working to relieve the pressure!

Medical Conditions

Acute Sinusitis

Addisions Disease

Adrenal Insufficiency

Aneamia

Behcets Disease

Blood Clotting Disorders

Chronic Ear Diseases

Chronic Kidney Disease

Gluten Intolerance

Guillian Barre Syndrome

Head Trauma

Hyperadrenalism

Hyperaldostronism

Hypercoagulable States

Hyperthyroidism

Hypoparathyroidism

Infections

Kidney Failure

Lactose Intolerance

Leukaemia

Lupus

Lyme Disease

Malnutrition

Menopause

Obstructive Sleep Apnoea

Overactive & Underactive Thyroid/Hypothyroidism & Hyperthyroidism

Polycystic Ovary Syndrome

Pregnancy

Protein Malnutrition

Sarcoidosis

SLE

Vitamin D Deficiency

Drugs

All trans retinoic acid (used in the treatment of Promyelocytic Leukaemia)

Amiodarone

Contraceptives

Corticosteroid treatment (including withdrawal)

Cyclosporine

Excessive ingestions of Vitamin A (Hypervitaminosis A)

Growth Hormone

Indomethicin

Isotretinoin (Accutane) for acne

Lithium

Minocycline

Nalidixic Acid

Nitrofurantoin

Phenytoin

Steroid withdrawal

Tetracycline

Thyroid Replacement

Vasopressin